



11175 Flintkote Ave., Suite E, San Diego, CA 92121
Tel: (858) 642-0978 Fax (858) 642-0937 Toll free: 1-888-723-GENE
E-mail: info@imgenex.com; web site: www.imgenex.com

Recombinant Ovine Growth Hormone (rOGH)

◀ **Description:**

Recombinant Ovine Growth Hormone produced in E.Coli is a single, non-glycosylated polypeptide chain containing 200 amino acids and having a molecular mass of 22015 Dalton.

The rOGH is purified by proprietary chromatographic techniques.

◀ **Source:**

Escherichia Coli.

◀ **Lot Number:**

◀ **Quantity Shipped:**

◀ **Physical Appearance:**

Sterile Filtered White lyophilized (freeze-dried) powder.

◀ **Formulation:**

The protein was lyophilized from a concentrated (1mg/ml) solution with 0.0045mM NaHCO₃.

◀ **Solubility:**

It is recommended to reconstitute the lyophilized rOGH in sterile 18MΩ-cm H₂O not less than 100μg/ml, which can then be further diluted to other aqueous solutions.

◀ **Stability:**

Lyophilized rOGH although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution rOGH should be stored at 4°C between 2-7 days and for future use below -18°C.

For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

Please prevent freeze-thaw cycles.

◀ **Purity:**

Greater than 98.0% as determined by:

- (a) Analysis by RP-HPLC.
- (b) Anion-exchange FPLC.
- (c) Analysis by reducing and non-reducing SDS-PAGE Silver Stained gel.

◀ **Amino acid sequence:**

The sequence of the first five N-terminal amino acids was determined and was found to be Met-Ala-Phe-Pro-Ala.

◀ **Dimers and aggregates:**

Less than 1% as determined by silver-stained SDS-PAGE gel analysis.

◀ **Biological Activity:**

ProSpec's rOGH is fully biologically active when compared to standard.

The activity as determined by the dose-dependant stimulation of the proliferation FDCP13B9 cells.

◀ **Endotoxin:**

Less than 0.1 ng/μg (IEU/μg) of rOGH.

◀ **Protein content:**

Protein quantitation was carried out by two independent methods:

1. UV spectroscopy at 280 nm using the absorbency value of 0.7 as the extinction coefficient for a 0.1% (1mg/ml) solution. This value is calculated by the PC GENE computer analysis program of protein sequences (IntelliGenetics).
2. Analysis by RP-HPLC, using a calibrated solution of rOGH as a Reference Standard.

◀ **Usage:**

This material is offered by ProSpec-TechnoGene for research, laboratory or further manufacturing purposes.